

Smooth globally hyperbolic splittings and temporal functions

Antonio N. Bernal and Miguel Sánchez

Depto. Geometría y Topología, Universidad de Granada,
Facultad de Ciencias, Avda. Fuentenueva s/n, E-18071 Granada, Spain

February 7, 2008

Abstract

Geroch's theorem about the splitting of globally hyperbolic spacetimes is a central result in global Lorentzian Geometry. Nevertheless, this result was obtained at a topological level, and the possibility to obtain a metric (or, at least, smooth) version has been controversial since its publication in 1970. In fact, this problem has remained open until a definitive proof, recently provided by the authors. Our purpose is to summarize the history of the problem, explain the smooth and metric splitting results (including smoothability of time functions in stably causal spacetimes), and sketch the ideas of the solution.

Keywords: Lorentzian manifold, globally hyperbolic, Cauchy hypersurface, smooth splitting, Geroch's theorem, stably causal spacetime, time function.

MSC: 53C50, 83C05

1 Introduction

Geroch's theorem [13] is a cut result in Lorentzian Geometry which, essentially, establishes the equivalence for a spacetime (M, g) between: (A) global hyperbolicity, i.e., strong causality plus the compactness of $J^+(p) \cap J^-(q)$ for all $p, q \in M$, and (B) the existence of a Cauchy hypersurface S , i.e. S is an achronal subset which is crossed exactly once by any inextendible timelike curve¹. Even more, the proof is carried out by finding two elements with interest in its own right: (1) an onto (global) time function $t : M \rightarrow \mathbb{R}$ (i.e. the onto function t is continuous and increases strictly on any causal curve) such that each level $t^{-1}(t_0)$, $t_0 \in \mathbb{R}$, is a Cauchy hypersurface and, then, (2) a global topological splitting $M \equiv \mathbb{R} \times S$ such that each slice $\{t_0\} \times S$ is a Cauchy hypersurface. Recall also that the existence of a time function t characterizes stably causal spacetimes

¹In particular, S is a topological hypersurface (without boundary), and it is also crossed at some point -perhaps even along a segment- by any lightlike curve.

(those causal spacetimes which remain causal under C^0 perturbations of the metric).

The possibility to smooth these topological results or continuous elements, have remained as an open folk question since its publication. In fact, Sachs and Wu claimed in their survey on General Relativity in 1977 [20, p. 1155]:

This is one of the folk theorems of the subject. It is not difficult to prove that every Cauchy surface is in fact a Lipschitzian hypersurface in M [19]. However, to our knowledge, an elegant proof that his Lipschitzian submanifold can be smoothed out [to such a smooth Cauchy hypersurface] is still missing.

Recall that here only the necessity to prove the smoothness of some Cauchy hypersurface S is claimed but, obviously, this would be regarded as a first step towards a fully satisfactory solution of the problem, among the following three:

- (i) To ensure the existence of a (smooth) *spacelike* S (necessarily, such an S will be crossed exactly once by any inextendible causal curve).
- (ii) To find not only a time function but also a “temporal” one, i.e., smooth with timelike gradient (even for any stably causal spacetime).
- (iii) To prove that any globally hyperbolic spacetime admits a smooth splitting $M = \mathbb{R} \times S$ with Cauchy hypersurfaces slices $\{t_0\} \times S$ orthogonal to ∇t (and, thus, with a metric without cross terms between $\mathbb{R}$ and S).

Among the concrete applications of (i), recall, for example: (a) Cauchy hypersurfaces are the natural regions to pose (smooth!) initial conditions for hyperbolic equations, as Einstein’s ones, or (b) differentiable achronal hypersurfaces (as those with prescribed mean curvature [9]) can be regarded as differentiable graphs on any smooth Cauchy hypersurface. The smoothness of a time function t claimed in (ii), would yield the possibility to use its gradient, which can be used to split any stably causal spacetime, as in [12]. The applications of the full smooth splitting result (iii) include topics such as Morse Theory for lightlike geodesics [22], quantization [10] or the possibility to use variational methods [15, Chapter 8]; it also opens the possibility to strengthen other topological splitting results [16] into smooth ones.

Recently, we have given a full solution to these three questions (i)—(iii) [3, 4]. Our purpose in this talk is, first, to summarize the history of the problem and previous attempts (Section 2) as well as the background results (Section 3). In the two following sections, our main results are stated and the ideas of the proofs sketched². Concretely, Section 4 is devoted to the construction of a smooth and spacelike S , following [3], and Section 5 to the full splitting of globally hyperbolic spacetimes, plus the existence of temporal functions in stably causal ones, following [4]. The reader is referred to the original references [3, 4] for detailed proofs and further discussions.

²The results of Section 5 has been obtained just some weeks ago, and they were not known at the time of the meeting in November 2003, but they are sketched here because of their obvious interest for our problem.

2 A brief history of time functions

As far as we know, the history of the smoothing splitting theorem can be summarized as follows.

1. Geroch published his result in 1970 [13], stating clearly all the results at a topological level. Penrose cites directly Geroch's paper in his book (1972), and regards explicitly the result as topological [19, Theorems 5.25, 5.26]. A subtle detail about his statement of the splitting result [19, Theorem 5.26] is the following. It is said there that, fixed $x \in S$, the curve $t \rightarrow \gamma(t) = (t, x) \in \mathbb{R} \times S$ is timelike. Recall that this curve does satisfy $t < t' \Rightarrow \gamma(t) << \gamma(t')$, but it is not necessarily smooth (its parameter is only continuous). In Penrose's definition [19, pp. 2, 3] timelike curves are assumed smooth, but later on [19, p. 17] the definition of causal curves is extended; in general, they will be just Lipschitzian, as previous curve γ .
2. Seifert's thesis (1968) develops a smoothing regularization procedure for time functions, which would yield the global splitting (Theorem 5.10 below). His smoothing results were published in 1977, [21], but the paper contains major gaps which could not be filled later.
3. In Hawking, Ellis' book (1973), the equivalence between stable causality and the existence of a (continuous) time function is achieved, by following a modification of Geroch's technique. Nevertheless, they assert [14, Proposition 6.4.9] that stable causality holds if and only if a (smooth) function with timelike gradient exists. Unfortunately, they refer for the details of the smoothing result to Seifert's thesis. Even more, in [14, Proposition 6.6.8], Geroch's result is stated at a topological level, but they refer to the possibility to smooth the result at the end of the proof. Nevertheless, again, the cited technique is the same for time functions.
4. In 1976, Budic and Sachs carried out a smoothing for *deterministic* spacetimes. One year later Sachs and Wu [20] posed the smoothing problem as a folk topic in General Relativity, in the above quoted paragraph.
5. The prestige and fast propagation of some of the previous references, made even the strongest splitting result be cited as proved in many references, including new influential references or books (for example, [10, 15, 22, 23]). But this is not the case for most references in pure Lorentzian Geometry, as O'Neill's book [18] (or, for example, [5, 9, 11, 16]). Even more, in Beem, Ehrlich's book (1981) Sachs and Wu's claim is referred explicitly [1, p. 31].
6. In 1988, Dieckmann claimed to prove the "folk question"; nevertheless, he cited Seifert's at the crucial step [7, Proof of Theorem 1]. More precisely, his study (see [8]) clarified other point in Geroch's proof, concerning the existence of an appropriate finite measure on the manifold. Even though the straightforward way to construct this measure in Hawking-Ellis' book

[14, proof of Proposition 6.4.9] is correct, neither these authors nor Geroch considered the necessary abstract properties that such a measure must fulfill (in particular, the measure of the boundaries $\partial I^+(p)$ must be 0). Under this approach, on one hand, the admissible measures for the proof of Geroch's theorem are characterized and, on the other, a striking relationship between continuity of volume functions and reflectivity is obtained.

In the 2nd edition of Beem-Ehrlich's book, in collaboration with Easley (1996), these improvements by Dieckmann are stressed, but Geroch's result is regarded as topological, and the reference to Sachs and Wu's claim is maintained [2, p. 65].

In general, continuous functions can be approximated by smooth functions. Thus, a natural way to proceed would be to approximate the continuous time function provided in Geroch's result, by a smooth one. Nevertheless, this intuitive idea has difficulties to be formalized. Thus, our approach has been different. First, we managed to smooth a Cauchy hypersurface [3] and, then, we constructed the full time function with the required properties [4].

3 Setup and previous results

Detailed proofs of the fact that the existence of a Cauchy hypersurface implies global hyperbolicity can be found, for example, in [13, 14, 18]. We will be interested in the converse, and then Geroch's results can be summarized in Theorem 3.2, plus Lemma 3.1 and Corollary 3.3.

Lemma 3.1 *Let M be a (C^k) -spacetime which admits a C^r -Cauchy hypersurface S , $r \in \{0, 1, \dots, k\}$. Then M is C^r -diffeomorphic to $\mathbb{R} \times S$ and all the C^r -Cauchy hypersurfaces are C^r diffeomorphic.*

This lemma is proven by moving S with the flow Φ_t of any complete timelike vector field. Thus, the (differentiable) hypersurfaces at constant $t \in \mathbb{R}$ are not necessarily Cauchy nor even spacelike, except for $t = 0$.

Theorem 3.2 *Assume that the spacetime M is globally hyperbolic. Then there exists a continuous and onto map $t : M \rightarrow \mathbb{R}$ satisfying:*

- (1) $S_a := t^{-1}(a)$ is a Cauchy hypersurface, for all $a \in \mathbb{R}$.
- (2) t is strictly increasing on any causal curve.

Function t is constructed as

$$t(z) = \ln (\text{vol}(J^-(z))/\text{vol}(J^+(z)))$$

for a (suitable) finite measure on M and, thus, global hyperbolicity implies just its continuity. Finally, combining both previous results,

Corollary 3.3 *Let M be a globally hyperbolic spacetime. Then there exists a homeomorphism*

$$\Psi : M \rightarrow \mathbb{R} \times S_0, \quad z \rightarrow (t(z), \rho(z)), \quad (1)$$

which satisfies:

(a) *Each level hypersurface $S_t = \{z \in M : t(z) = t\}$ is a Cauchy hypersurface.*

(b) *Let $\gamma_x : \mathbb{R} \rightarrow M$ be the curve in M characterized by:*

$$\Psi(\gamma_x(t)) = (t, x), \quad \forall t \in \mathbb{R}.$$

Then the continuous curve γ_x is timelike in the following sense:

$$t < t' \Rightarrow \gamma_x(t) << \gamma_x(t').$$

Remark 3.4 If function t in Corollary 3.3 were smooth with timelike gradient, then the spacetime (M, g) would be isometric to $\mathbb{R} \times \mathcal{S}, \langle \cdot, \cdot \rangle = -\beta dt^2 + \bar{g}$, where $\bar{g}$ is a (positive definite) Riemannian metric on each slice $t = \text{constant}$. The splitting is then obtained projecting M to a fixed level hypersurface by means of the flow of ∇t . In what follows, a smooth function $\mathcal{T}$ with past-pointing timelike gradient $\nabla \mathcal{T}$ will be called *temporal* (and it is necessarily a time function).

For the smoothing procedure, some properties of Cauchy hypersurfaces will be needed. Concretely, by using a result on intersection theory, the following one can be proven [3, Section 3], [11, Corollary 2]:

Proposition 3.5 *Let S_1, S_2 be two Cauchy hypersurfaces of a globally hyperbolic spacetime with $S_1 << S_2$ (i.e., $S_2 \subset I^+(S_1)$), and S be a connected closed spacelike hypersurface (without boundary):*

(A) *If $S_1 << S$ then S is achronal, and a graph on all S_1 for the decompositions in Corollaries 3.1, 3.3.*

(B) *If $S_1 << S << S_2$ then S is a Cauchy hypersurface.*

4 Smooth spacelike Cauchy hyperfurfaces

In this section, we sketch the proof of:

Theorem 4.6 *Any globally hyperbolic spacetime admits a smooth spacelike Cauchy hypersurface S .*

(In what follows, “smooth” will mean with the same order of differentiability of the spacetime).

From Proposition 3.5, given two Cauchy hypersurfaces $S_1 << S_2$ as in Theorem 3.2 (with $S_{t_i} \equiv S_i$; $t_1 < t_2$), it is enough to construct a connected closed spacelike hypersurface S with $S_1 << S << S_2$. And, in order to prove this, it suffices:

Proposition 4.7 *Let M be a globally hyperbolic spacetime with topological splitting $\mathbb{R} \times S$ as in Corollary 3.3, and fix $S_1 \ll S_2$. Then there exists a smooth function*

$$h : M \rightarrow [0, \infty)$$

which satisfies:

1. $h(t, x) = 0$ if $t \leq t_1$.
2. $h(t, x) > 1/2$ if $t = t_2$.
3. *The gradient of h is timelike and past-pointing on the open subset $V = h^{-1}((0, 1/2)) \cap I^-(S_2)$.*

In fact, recall that, given such a function h , each $s \in (0, 1/2)$ is a regular value of the restriction of h to $J^-(S_2)$. Thus, $S_s^h := h^{-1}(s) \cap J^-(S_2)$ is a closed smooth spacelike hypersurface with $S_1 \ll S_s^h \ll S_2$ and, then, a Cauchy hypersurface (in principle, Proposition 3.5(B) can be applied to any connected component of S_s^h , but one can check that, indeed, S_s^h is connected).

The construction of function h is carried out in two closely related steps. The first one is a local step: to construct, around each $p \in S_2$, a function h_p with the suitable properties, stated in Lemma 4.8. The second step is to construct (global) function h from the h_p 's. This function will be constructed directly as a sum³ $h = \sum h_i$ for suitable $h_i \equiv h_{p_i}$. This fact must be taken into account for the properties of the h_p 's in the first step:

Lemma 4.8 *Fix $p \in S_2$, and a convex neighborhood of p , $\mathcal{C}_p \subset I^+(S_1)$ (that is, $\mathcal{C}_p$ is a normal starshaped neighborhood of any of its points).*

Then there exists a smooth function

$$h_p : M \rightarrow [0, \infty)$$

which satisfies:

- (i) $h_p(p) = 1$.
- (ii) *The support of h_p (i.e., the closure of $h_p^{-1}(0, \infty)$) is compact and included in $\mathcal{C}_p \cap I^+(S_1)$.*
- (iii) *If $q \in J^-(S_2)$ and $h_p(q) \neq 0$ then $\nabla h_p(q)$ is timelike and past-pointing.*

Sketch of proof. Function h_p is taken in a neighbourhood of $\mathcal{C}_p \cap J^-(S_2)$ as:

$$h_p(q) = e^{d(p', p)^{-2}} \cdot e^{-d(p', q)^{-2}},$$

where d is the time-separation (Lorentzian distance) on $\mathcal{C}_p$, and p' is a fixed suitably chosen point in the past of p .

³In Riemannian Geometry, global objects are constructed frequently from local ones by using partitions of the unity. Nevertheless, the causal character of the gradient of functions in the partition are, in principle, uncontrolled. Then, the underlying idea to construct h is to use the paracompactness of M (which is implied by the existence of a Lorentzian –or semi-Riemannian– metric) avoiding to use a partition of the unity.

Now, the second step is carried out by taking advantage directly of the paracompactness of the manifold. Concretely, function $h = \sum_i h_i$ is obtained by choosing the h_i 's from the following lemma, with the W_α 's equal to $h_p^{-1}(1/2, \infty)$, and $p \in S_2$ (see [3] for details):

Lemma 4.9 *Let d_R be the distance on M associated to any auxiliary complete Riemannian metric g_R . Let S_2 be a closed subset of M and $\mathcal{W} = \{W_\alpha, \alpha \in \mathcal{I}\}$ a collection of open subsets of M which cover S_2 . Assume that each W_α is included in an open subset $\mathcal{C}_\alpha$ and the d_R -diameter of each $\mathcal{C}_\alpha$ is smaller than 1. Then there exist a subcollection $\mathcal{W}' = \{W_j : j \in \mathbb{N}\} \subset \mathcal{W}$ which covers S_2 and is locally finite (i.e., for each $p \in \cup_j W_j$ there exists a neighborhood V such that $V \cap W_j = \emptyset$ for all j but a finite set of indexes). Moreover, the collection $\{\mathcal{C}_j : j \in \mathbb{N}\}$ (where each $W_j \in \mathcal{W}'$ is included in the corresponding $\mathcal{C}_j$) is locally finite too.*

5 Temporal functions and the full splitting

Now, our aim is to sketch the proof of the following theorem.

Theorem 5.10 *Let (M, g) be a globally hyperbolic spacetime. Then, it is isometric to the smooth product manifold*

$$\mathbb{R} \times \mathcal{S}, \quad \langle \cdot, \cdot \rangle = -\beta d\mathcal{T}^2 + \bar{g}$$

where $\mathcal{S}$ is a smooth spacelike Cauchy hypersurface, $\mathcal{T} : \mathbb{R} \times \mathcal{S} \rightarrow \mathbb{R}$ is the natural projection, $\beta : \mathbb{R} \times \mathcal{S} \rightarrow (0, \infty)$ a smooth function, and $\bar{g}$ a 2-covariant symmetric tensor field on $\mathbb{R} \times \mathcal{S}$, satisfying:

1. $\nabla \mathcal{T}$ is timelike and past-pointing on all M , that is, function $\mathcal{T}$ is temporal.
2. Each hypersurface $\mathcal{S}_\mathcal{T}$ at constant $\mathcal{T}$ is a Cauchy hypersurface, and the restriction $\bar{g}_\mathcal{T}$ of $\bar{g}$ to such a $\mathcal{S}_\mathcal{T}$ is a Riemannian metric (i.e. $\mathcal{S}_\mathcal{T}$ is spacelike).
3. The radical of $\bar{g}$ at each $w \in \mathbb{R} \times \mathcal{S}$ is $\text{Span} \nabla \mathcal{T}$ ($= \text{Span } \partial_\mathcal{T}$) at w .

Essentially, it is enough for the proof to obtain a temporal function $\mathcal{T} : M \rightarrow \mathbb{R}$ such that each level hypersurface is Cauchy, see Remark 3.4. The existence of such a $\mathcal{T}$ is carried out in three steps.

Step 1: time step functions would solve the problem. Let $t \equiv t(z)$ be a continuous time function as in Geroch's Theorem 3.2. Fixed $t_- < t \in \mathbb{R}$, we have proven in Section 4 the existence of a smooth Cauchy hypersurface $\mathcal{S}$ contained in $t^{-1}(t_-, t)$; this hypersurface is obtained as the regular value of certain function $h \equiv h_t$ with timelike gradient on $t^{-1}(t_-, t]$. As t_- approaches t , $\mathcal{S}$ can be seen as a smoothing of S_t ; nevertheless $\mathcal{S}$ always lies in $I^-(S_t)$. Now, we claim that the required splitting of the spacetime would be obtained if we could strengthen the

requirements on this function h_t , ensuring the existence of a *time step function* τ_t around each S_t . Essentially such a τ_t is a function with timelike gradient on a neighborhood of S_t (and 0 outside) with level Cauchy hypersurfaces which cover a rectangular neighbourhood of S_t :

Lemma 5.11 *All the conclusions of Theorem 5.10 will hold if the globally hyperbolic spacetime M admits, around each Cauchy hypersurface S_t , $t \in \mathbb{R}$, a (time step) function $\tau_t : M \rightarrow \mathbb{R}$ which satisfies:*

1. $\nabla \tau_t$ is timelike and past-pointing where it does not vanish, that is, in the interior of its support $V_t := \text{Int}(\text{Supp}(\nabla \tau_t))$.
2. $-1 \leq \tau_t \leq 1$.
3. $\tau_t(J^+(S_{t+2})) \equiv 1$, $\tau_t(J^-(S_{t-2})) \equiv -1$.
4. $S_{t'} \subset V_t$, for all $t' \in (t-1, t+1)$; that is, the gradient of τ_t does not vanish in the rectangular neighborhood of S , $t^{-1}(t-1, t+1) \equiv (t-1, t+1) \times S$.

Sketch of proof. Consider such a function τ_k for $k \in \mathbb{Z}$, and define the (locally finite) sum $\mathcal{T} = \tau_0 + \sum_{k=1}^{\infty} (\tau_{-k} + \tau_k)$. One can check that $\mathcal{T}$ fulfills the required properties in Remark 3.4, in fact: (a) $\mathcal{T}$ is temporal because subsets $V_{t=k}$, $k \in \mathbb{Z}$ cover all M (and the timelike cones are convex), and (b) the level hypersurfaces of $\mathcal{T}$ are Cauchy because, for each inextendible timelike curve $\gamma : \mathbb{R} \rightarrow M$ parameterized with $\mathcal{T}$, $\lim_{s \rightarrow \pm\infty} (\mathcal{T}(\gamma(s))) = \pm\infty$.

■

Step 2: constructing a weakening of a time step function. Lemma 5.11 reduces the problem to the construction of a time step function τ_t for each t . We will start by constructing a function $\hat{\tau}_t$ which satisfies all the conditions in that lemma but the last one, which is replaced by:

4. $S_t \subset V_t$.

The idea for the construction of such a $\hat{\tau}_t$ is the following. Consider function h in Lemma 4.7 for $t_1 = t-1, t_2 = t$. From its explicit construction, it is straightforward to check that h can be also assumed to satisfy: ∇h is timelike and past-pointing on a neighborhood $U' \subset I^-(S_{t+1})$ of S_t . Thus, putting $U = U' \cup I^-(S_t)$ (U satisfies $J^-(S_{t-1}) \subset U \subset I^-(S_{t+1})$), we find a function $h^+ : M \rightarrow \mathbb{R}$ which satisfies:

- (i⁺) $h^+ \geq 0$ on U , with $h^+ \equiv 0$ on $I^-(S_{t-1})$.
- (ii⁺) If $p \in U$ with $h^+(p) > 0$ then $\nabla h^+(p)$ is timelike and past-pointing.
- (iii⁺) $h^+ > 1/2$ (and, thus, its gradient is timelike past-pointing) on $J^+(S_t) \cap U$.

Even more, a similar reasoning yields a function $h^- : M \rightarrow \mathbb{R}$ for this same U which satisfies:

- (i⁻) $h^- \leq 0$, with $h^- \equiv 0$ on $M \setminus U$.
- (ii⁻) If $\nabla h^-(p) \neq 0$ at $p (\in U)$ then $\nabla h^-(p)$ is timelike past-pointing.
- (iii⁻) $h^- \equiv -1$ on $J^-(S_t)$.

Now, as $h^+ - h^- > 0$ on all U , we can define:

$$\hat{\tau}_t = 2 \frac{h^+}{h^+ - h^-} - 1$$

on U , and constantly equal to 1 on $M \setminus U$. A simple computation shows that $\nabla \hat{\tau}_t$ does not vanish wherever either $h^- \nabla h^+$ or $h^+ \nabla h^-$ does not vanish (in particular, on S_t) and, then, it fulfills all the required conditions.

Step 3: construction of a true time step function. Now, our aim is to obtain a function $\tau (\equiv \tau_t)$ which satisfies not only the requirements of previous $\hat{\tau} \equiv \hat{\tau}_t$ but also the stronger condition 4 in Lemma 5.11. Fix any compact subset $K \subset t^{-1}([t-1, t+1])$. From the construction of $\hat{\tau}$, it is easy to check that $\hat{\tau}$ can be chosen with $\nabla \hat{\tau}$ non-vanishing on K . Now, choose a sequence $\{G_j : j \in \mathbb{N}\}$ of open subsets such that:

$$\overline{G_j} \text{ is compact, } \overline{G_j} \subset G_{j+1} \quad M = \cup_{j=1}^{\infty} G_j,$$

and let $\hat{\tau}[j]$ be the corresponding sequence of functions type $\hat{\tau}$ with gradients non-vanishing on:

$$K_j = \overline{G_j} \cap t^{-1}([t-1, t+1]).$$

Essentially, the required time step function is:

$$\tau = \sum_{j=1}^{\infty} \frac{1}{2^j A_j} \hat{\tau}[j],$$

where each A_j is chosen to make τ smooth (fixed a locally finite atlas on M , each A_j bounds on $\overline{G_j}$ each function $\hat{\tau}[j]$ and its partial derivatives up to order j in the charts of the atlas which intersect G_j). Then, the gradient of τ is timelike wherever one of the gradients $\nabla \hat{\tau}[j]$ does not vanish (in particular, on $t^{-1}([t-1, t+1])$). Moreover, τ is equal to constants (which can be rescaled to ± 1) on $t^{-1}((-\infty, t-2]), t^{-1}([t+2, \infty))$, as required. ■

Finally, it is worth pointing out that similar arguments work to find a smooth time function on any spacetime (even non-globally hyperbolic) which admits a continuous time function t .

Theorem 5.12 *Any spacetime M which admits a (continuous) time function (i.e., is stably causal) also admits a temporal function $\mathcal{T}$.*

Sketch of proof. Notice that each level continuous hypersurface S_t is a Cauchy hypersurface in its Cauchy development $D(S_t)$. Moreover, any time step function τ_t around S_t in $D(S_t)$ can be extended to all M (making τ_t equal to 1 on $I^+(S_t) \cap (M \setminus D(S))$, and to -1 on $I^-(S_t) \cap (M \setminus D(S))$). Then, sum suitable time step functions as in Step 3 above.

■

Acknowledgments

The second-named author has been partially supported by a MCyT-FEDER Grant BFM2001-2871-C04-01.

References

- [1] J.K. BEEM AND P.E. EHRLICH, *Global Lorentzian geometry*, Monographs and Textbooks in Pure and Applied Math., **67**, Marcel Dekker, Inc., New York (1981).
- [2] J.K. BEEM, P.E. EHRLICH AND K.L. EASLEY, *Global Lorentzian geometry*, Monographs Textbooks Pure Appl. Math. **202**, Dekker Inc., New York (1996).
- [3] A. N. BERNAL AND M. SÁNCHEZ, On Smooth Cauchy Hypersurfaces and Geroch's Splitting Theorem, *Commun. Math. Phys.*, **243** (2003) 461-470.
- [4] A. N. BERNAL AND M. SÁNCHEZ, Smoothness of time functions and the metric splitting of globally hyperbolic spacetimes, preprint, gr-qc/0401112
- [5] R. BUDIC, J. ISENBERG, L. LINDBLOM AND P.B. YASSKIN, On determination of Cauchy surfaces from intrinsic properties. *Comm. Math. Phys.* **61** (1978), no. 1, 87–95.
- [6] R. BUDIC AND R.K. SACHS, Scalar time functions: differentiability. Differential geometry and relativity, pp. 215–224, *Mathematical Phys. and Appl. Math.*, Vol. 3, Reidel, Dordrecht, 1976.
- [7] J. DIECKMANN, Volume functions in general relativity, *Gen. Relativity Gravitation* **20** (1988), no. 9, 859–867.
- [8] J. DIECKMANN, Cauchy surfaces in a globally hyperbolic space-time, *J. Math. Phys.* **29** (1988), no. 3, 578–579.
- [9] K. ECKER AND G. HUISKEN, Parabolic methods for the construction of spacelike slices of prescribed mean curvature in cosmological spacetimes, *Comm. Math. Phys.* **135** (1991), no. 3, 595–613.
- [10] E.P. FURLANI, Quantization of massive vector fields in curved space-time, *J. Math. Phys.* **40** (1999), no. 6, 2611–2626.
- [11] G.J. GALLOWAY, Some results on Cauchy surface criteria in Lorentzian geometry, *Illinois J. Math.*, **29** (1985), no. 1, 1–10.
- [12] E. GARCA-RO AND D.N. KUPELI, Some splitting theorems for stably causal spacetimes. *Gen. Relativity Gravitation* **30** (1998), no. 1, 35–44.
- [13] R. GEROCH, Domain of dependence, *J. Math. Phys.* **11** (1970) 437–449.

- [14] S.W. HAWKING AND G.F.R. ELLIS, *The large scale structure of space-time*, Cambridge Monographs on Mathematical Physics, No. 1. Cambridge University Press, London-New York (1973).
- [15] A. MASIELLO, *Variational methods in Lorentzian geometry*, Pitman Res. Notes Math. Ser. **309**, Longman Sci. Tech., Harlow (1994).
- [16] R.P.A.C. NEWMAN, The global structure of simple space-times, *Comm. Math. Phys.* **123** (1989), no. 1, 17–52.
- [17] K. NOMIZU AND H. OZEKI, The existence of complete Riemannian metrics *Proc. Amer. Math. Soc.* **12** (1961) 889–891.
- [18] B. O’NEILL, *Semi-Riemannian geometry with applications to Relativity*, Academic Press Inc., New York (1983).
- [19] R. PENROSE, *Techniques of Differential Topology in Relativity*, CBMS-NSF Regional Conference Series in applied Mathematics, SIAM, Philadelphia (1972).
- [20] R.K. SACHS AND H. WU, General relativity and cosmology, *Bull. Amer. Math. Soc.* **83** (1977), no. 6, 1101–1164.
- [21] H.-J. SEIFERT, Smoothing and extending cosmic time functions, *Gen. Relativity Gravitation* **8** (1977), no. 10, 815–831.
- [22] K. UHLENBECK, A Morse theory for geodesics on a Lorentz manifold. *Topology* **14** (1975), 69–90.
- [23] R.M. WALD, *General Relativity*, Univ. Chicago Press, 1984.